

NDSU Carrington Research Extension Center
2010 Variety Trial Data

Emmer - Organic	Carrington
------------------------	-------------------

Variety	Days to Heading	Plant Lodge 0-9	Plant Height inch	1000 KWT gram	Test Weight lb/bu	----- Seed Yield -----	
						2010 lb/ac	2-yr. Avg.
Bowman	67.3	0.8	35.4	49.0	36.0	1845.1	1684.9
Common-H	67.5	0.8	33.5	46.0	35.5	1747.4	1801.7
Common-M	66.5	0.3	36.2	54.5	37.1	1931.9	1832.0
Common-R	66.8	0.3	35.6	52.3	37.4	1601.2	1733.0
Common-MC	67.3	0.5	34.3	51.4	36.2	2014.3	2031.4
Debra	62.5	0.0	23.4	36.0	55.0	1367.5	1295.4
Lucille	66.8	0.5	37.8	64.0	35.7	2282.1	2303.1
ND Common	67.0	1.3	35.0	57.0	37.9	2309.9	2248.2
PI254148	60.2	0.4	22.4	55.9	31.3	1524.2	--
Vernal	66.8	0.5	37.4	64.6	37.5	1940.2	1887.9
Red Vernal	67.3	1.3	36.4	60.2	35.5	2100.8	2158.8
Mean	66.1	0.6	33.6	53.6	37.9	1865.7	--
C.V.%	1.4	122.0	7.0	9.3	5.6	19.8	--
LSD.05	1.3	1.0	3.4	7.2	3.0	542.7	--

Planting Date = April 22; Harvest Date = August 2; Previous Crop = Cover crop (winter rye, annual ryegrass, hairy vetch, berseem clover, turnip, and rape)

*Trial was infected with wheat streak mosaic from volunteer winter wheat.